

September 1996

HUTP-96/A044

IFP-735-UNC

Cabibbo Mixing and the Search for CP Violation

Paul H. Frampton^(a) and Sheldon L. Glashow^(b)

^(a) *University of North Carolina, Chapel Hill, NC 27599-3255*

^(b) *Harvard University, Cambridge, MA 02138.*

Abstract

We examine certain extensions of the standard model in which CP violation is spontaneous and the strong CP problem is resolved. In these models, the 3×3 quark mixing matrix is neither real nor unitary. However, to a precision of 0.1%, it is real and orthogonal. There are no readily observable CP -violating effects besides those in the neutral kaon system.

Typeset using REVTeX

The Standard Model is often regarded as a correct and complete description of all elementary-particle phenomena, but its depiction of CP violation is uncertain. Although the effect was first observed three decades ago [1], only one CP -violating parameter (ϵ_K for neutral kaons) is precisely measured. Others are indecisively measured (*e.g.*, ϵ'/ϵ) or merely constrained (*e.g.*, the electric dipole moment of the neutron). More data is essential to achieve a definitive theory of CP violation, such as forthcoming studies of neutral B decays that will contribute to the evaluation of the parameters of quark mixing and provide important tests of the standard model [2–4]. However, the strong CP problem cannot be solved in the context of the standard model. Put in a nutshell, there is no natural way to control the phase of the determinant of the quark mass matrix without imposing CP conservation on the Lagrangian.

In this paper, we explore CP -violation in the context of an ‘Aspon model’ [5–8] wherein the violation of CP is spontaneous. In particular, we add to the 3-family standard model a vector-like $SU(2)$ doublet of quarks $(U, D)_L \oplus (U, D)_R$ together with two complex $SU(2)$ singlet scalars $\chi_{1,2}$.¹ Under an additional gauged $U(1)'$, these four representations carry the quantum numbers $X = +1, -1, +1, +1$, respectively. All conventional fields carry $X = 0$. CP invariance of the Lagrangian is assumed. Thus the Yukawa couplings of χ_i and of conventional Higg(s) fields are real. Spontaneous CP violation arises from complex vacuum expectation values acquired by the scalar singlets: $\langle \chi_1 \rangle = \kappa_1$ and $\langle \chi_2 \rangle = \kappa_2$.

In our scheme, quark masses are described by 4×4 matrices $\mathcal{M}_{u,d}$ taking the form:

$$\mathcal{M}_{u,d} = \begin{pmatrix} m_{u,d} & F \\ 0 & M \end{pmatrix}, \quad (1)$$

where $m_{u,d}$ are *real* 3×3 matrices resulting from the conventional Higgs mechanism and M is the Dirac mass of the added quark doublet. CP violation results from the complex

¹What we say is applicable to an alternative Aspon model where the added quarks are $SU(2)$ singlets.

off-diagonal matrix elements F that form a column vector. Its components:

$$F_k = h_k^1 \kappa_1 + h_k^2 \kappa_2. \quad (2)$$

involve the complex VEV's κ of the χ fields and their real Yukawa couplings:

$$L_{\text{Yukawa}} = \sum_{k=1}^3 \left(h_k^1 \chi_1 + h_k^2 \chi_2 \right) \left(\bar{u}_L^k U_R + \bar{d}_L^k D_R \right) + h.c. \quad (3)$$

We define the parameters x_k (assumed to be small) to be $x_k = F_k/M$.

The up and down mass matrices are diagonalized by bi-unitary transformations:

$$J_L^\dagger \mathcal{M}_u J_R = \text{diag}(m_u, m_c, m_t, M'), \quad (4)$$

$$K_L^\dagger \mathcal{M}_d K_R = \text{diag}(m_d, m_s, m_b, M''), \quad (5)$$

where we hereafter ignore the small differences between M , M' and M'' . The quark flavor mixing matrix is $C = J_L^\dagger K_L$, or more explicitly, $C_{\mu\nu} = J_{L\lambda\mu}^* K_{L\lambda\nu}$, where indices 1,2,3 (henceforth Latin) refer to known quarks and subscript 4 refers to the added heavy quarks. The CP -violating phases in C_{ij} are suppressed by the small parameters x_k^2 (the subscript is hereafter dropped).

A lower bound on x^2 arises from the need to describe observed CP violation in the neutral kaon system. Using Eq.(35) of [6] and requiring that $U(1)'$ break down above the electroweak scale (*i.e.*, $\kappa \geq 250$ GeV), we find $x^2 \geq 3 \times 10^{-5}$. With x^2 this large, box diagrams with χ exchange (in addition to conventional box diagrams) can contribute significantly to the neutral kaon mass difference. Unlike a superweak model [9], our description of neutral kaons yields a non-zero value of $|\epsilon'/\epsilon|$ bounded above by 2×10^{-3} [6].

Our model exhibits no strong CP problem in tree approximation. The determinants of $\mathcal{M}_{u,d}$ as given by (1) are real, so that $\bar{\theta} = 0$ at this order. An upper limit on x^2 is obtained from an examination of one-loop corrections to $\mathcal{M}_{u,d}$. For each of these matrices, we have

$$\delta \mathcal{M} = \begin{pmatrix} \delta m & \delta F \\ \delta G & \delta M \end{pmatrix}, \quad (6)$$

from which we find:

$$\begin{aligned}
\bar{\theta} &= \text{Arg Det}(\mathcal{M} + \delta\mathcal{M}) \\
&= \text{Im Tr}(\ln \mathcal{M}(1 + \mathcal{M}^{-1}\delta\mathcal{M})) \\
&\simeq \text{Im Tr} \mathcal{M}^{-1}\delta\mathcal{M}.
\end{aligned} \tag{7}$$

The last equality in (7) is perturbative in the one-loop radiative corrections. From (6) and (7) we deduce a useful expression for the one-loop correction to either of the phases $\bar{\theta}_{u,d}$:

$$\bar{\theta} \simeq \text{Im Tr} \left(m^{-1}\delta m + m^{-1}FM^{-1}\delta G + M^{-1}\delta M \right) \tag{8}$$

An explicit examination of the several bracketed terms in (8) reveals that all are real except the first. The term $m^{-1}\delta m$ yields a one-loop contribution to $\bar{\theta} = \bar{\theta}_u + \bar{\theta}_d$ of order $\lambda x^2/16\pi^2$, where λ is the real coefficient of the Higgs interaction $\lambda|\phi|^2|\chi|^2$. From $M_\phi^2 > (60 \text{ GeV})^2$ and the constraint $|\langle\chi\rangle| < 2 \text{ TeV}$ [6], we find $\lambda < 10^{-3}$. With x^2 at its lower limit, λ must be thirty times smaller than this upper bound to avoid a strong CP problem. It would be unnatural to set $\lambda < 10^{-5}$. Thus, we find as a plausible range for x^2 :

$$10^{-3} > x^2 > 3 \times 10^{-5}. \tag{9}$$

In an Aspon model, the quark mixing matrix is a complex unitary 4×4 matrix $C_{\mu\nu}$. Mixings of the conventional six quarks with one another are specified by the 3×3 matrix $C_{i,j}$, whose indices run from 1 to 3. C_{ij} is neither real nor unitary because of χ -induced mixings to the undiscovered quark doublet. However, these terms are $\sim x^2$. Thus C_{ij} , to a precision of at least 0.1%, is a real orthogonal matrix. It is a generalized Cabibbo matrix rather than a Kobayashi-Maskawa matrix. This is unfortunate for the search for CP violation in the beauty sector, but has other observable consequences.

In the standard model, the Kobayashi-Maskawa matrix V is complex and unitary. The sides of the “unitarity triangle” are unity and:

$$R_b = \left| \frac{V_{ud}V_{ub}}{V_{cd}V_{cb}} \right|, \quad R_t = \left| \frac{V_{td}V_{tb}}{V_{cd}V_{cb}} \right|. \tag{10}$$

The value of R_b has been measured. According to a recent review by M. Neubert [10], $R_b = 0.35 \pm 0.09$. The value of R_t cannot be extracted from experimental data alone.

Appeal must be made to a theoretical evaluation of the neutral B -meson mass difference using the standard model. Neubert's analysis yields $R_t = 0.99 \pm 0.22$. These results suggest a rather large value of the CP -violating angle β , namely $0.34 \leq \sin \beta \leq 0.75$.

In Aspon models, the matrix C_{ij} is orthogonal up to terms of order x^2 arising from mixings with unobserved quarks. Thus, we anticipate no readily observable manifestations of CP violation in the beauty sector. Furthermore, the unitarity triangle must degenerate into a straight line: $|R_b \pm R_t| = 1$. In this case, we cannot appeal to a theoretical calculation of the neutral B -meson mass difference because it depends on unknown parameters. On the other hand, the matrix C_{ij} with the neglect of terms $\sim x^2$ involves only three parameters. From data in hand, in this context, we obtain $R_t = 1 - \rho$ in the Wolfenstein parametrization [11], whence $R_t = 0.637 \pm 0.09$. We note that present data yields $R_b + R_t = 0.99 \pm 0.13$. This result is compatible with an approximately orthogonal mixing matrix and hence with our model.

This work was supported in part by the U. S. Department of Energy under Grant No. DE-FG05-85ER-40219, Task B, and by the National Science Foundation under Grant No. PHY-92-18167.

REFERENCES

- [1] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turlay, Phys. Rev. Lett. **13**, 138 (1964).
- [2] I. I. Bigi and A. I. Sanda, Nucl. Phys. **B193**, 85 (1981).
- [3] I. I. Bigi and A. I. Sanda, Nucl. Phys. **B281**, 41 (1987)
- [4] Y. Nir and H. R. Quinn, Ann. Rev. Nucl. Part. Sci. **42**, 211 (1992)
- [5] P. H. Frampton and T. W. Kephart, Phys. Rev. Lett. **66**, 1666 (1991).
- [6] P. H. Frampton and D. Ng, Phys. Rev. **D43**, 3034 (1991).
- [7] P. H. Frampton, T. W. Kephart, D. Ng and T. J. Weiler, Phys. Rev. Lett. **68**, 2129 (1992).
- [8] A. W. Ackley, P. H. Frampton, B. Kayser and C. B. Leung, Phys. Rev. **D50**, 3560 (1994)
- [9] L. Wolfenstein, Phys. Rev. Lett. **13**, 562 (1964).
- [10] M. Neubert, Int. J. Mod. Phys. **A11**, 4173 (1996).
- [11] L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1983).